

	LESSON PLAN	Semester : IV
	Sub Code & Name: MA 6452 STATISTICS AND NUMERICAL METHODS Unit: I Branch: MECH	

UNIT I TESTING OF HYPOTHESIS

Large sample test based on Normal distribution for single mean and difference of means - Tests based on t , Chi^2 and F distributions for testing means and variances - Contingency table (Test for Independency) - Goodness of fit.

Reference: Grewal. B.S., and Grewal. J.S., "Numerical Methods in Engineering and Science", 9th Edition, Khanna Publishers, New Delhi, 2007.

PART - A

Q.No.		BT Level	Competence	PO
1.	What is statistical hypothesis?(Nov/Dec-2017)	BTL -1	Remembering	PO1
2.	Define chi-square.?(Nov/Dec-2017)	BTL -1	Remembering	PO1
3.	Write type I and type II errors .(Apr/May-Nov/Dec-2013)(May/Jun-2016)	BTL -1	Remembering	PO1
4.	What are the assumptions in 't' distribution?(Nov/Dec-2016)(Apr/May-2015)	BTL -1	Remembering	PO1
5.	State the important properties of the t-distribution. (Apr/May-2015)	BTL -1	Remembering	PO1
6.	Write any three applications of Chi-Square distribution.(May/Jun-2014)	BTL -1	Remembering	PO1
7.	Define null and alternative hypothesis.	BTL -2	Understanding	PO2
8.	When do we use the t-distribution? (Nov/Dec-2016)	BTL -2	Understanding	PO2
9.	What is meant by level of significance? (Apr/May-2016)	BTL -2	Understanding	PO2
10.	Define Standard error and Critical region. (Nov/Dec-2016)	BTL -2	Understanding	PO2
11.	Write any two applications of 't'-distribution. (Nov/Dec-	BTL -3	Applying	PO3
12.	Write the condition for the application of χ^2 test.	BTL -3	Applying	PO3
13.	Write any three applications of 'F' distribution. (Nov/Dec-2015)	BTL -6	Creating	PO1,PO2, PO5
14.	State the important properties of F-distribution. (Nov/Dec-2011)	BTL -4	Analyzing	PO1,PO2, PO5
15.	Define sampling distribution. (Apr/May-2013)	BTL -4	Analyzing	PO1,PO2, PO5
16.	Define Chi-square test of goodness of fit. (Apr/May-2014)	BTL -3	Applying	PO5

17	Write down the form of the 95% confidence interval for the population mean in terms of population S.D.	BTL -5	Evaluating	PO1,PO2, PO5				
18.	What is the Standard error of the difference between the means of two large samples drawn from different populations with known SD's	BTL -5	Evaluating	PO1,PO2, PO5				
19.	What is the test statistic used to test the significance of the difference between small sample,mean and population?	BTL -6	Creating	PO1,PO2, PO5				
20.	What is the test statistic used to test the significance of the difference between the means of two small samples?	BTL -4	Analyzing	PO1,PO2, PO5				
21	Write down the formula of test statistic 'Z' to test the significance of difference between the means (large samples)	BTL -3	Applying	PO5				
22	Write down the formula of test statistic 'Z' to test the significance of difference between the proportions(large samples).	BTL -3	Applying	PO5				
23	What is the test statistic used to test the significance of the difference between the means of two small samples of same size,when the sample items are correlated?	BTL -6	Creating	PO1,PO2, PO5				
24	What are the expected frequency of 2x2 contingency table given below. <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>a</td> <td>b</td> </tr> <tr> <td>C</td> <td>d</td> </tr> </table>	a	b	C	d	BTL -4	Analyzing	PO1,PO2, PO12
a	b							
C	d							
25	Write down the 1% and 5% critical values for right tailed two tailed Tests.	BTL -4	Analyzing	PO1,PO2, PO12				
26	What is the difference between confidence limits and tolerance limits?	BTL -3	Applying	PO12				
27	What are the assumptions of large sample?	BTL -5	Evaluating	PO1,PO2, PO12				
28	What is test of goodness of fit?	BTL -3	Applying	PO12				
29	Define hypothesis	BTL -6	Creating	PO1,PO2, PO5				
30	What is meant by population?	BTL -4	Analyzing	PO1,PO2, PO5				
PART - B								

1.(a)	The mean life time of a sample of 100 light tubes produced by a company is found to be 1580 hours with standard deviation of 90 hours. Examine the hypothesis that the mean life time of the tubes produced by the company is 1600 hours. (Apr/May-2016)	BTL -1	Remembering	PO1																														
1. (b)	<p>Test of fidelity and selectivity of 190 radio receivers produced the results shown in the following table</p> <table border="1"> <thead> <tr> <th colspan="4">Fidelity</th> </tr> <tr> <th>Selectivity</th> <th>Low</th> <th>Average</th> <th>High</th> </tr> </thead> <tbody> <tr> <td>Low</td> <td>6</td> <td>12</td> <td>32</td> </tr> <tr> <td>Average</td> <td>33</td> <td>61</td> <td>18</td> </tr> <tr> <td>High</td> <td>13</td> <td>15</td> <td>0</td> </tr> </tbody> </table> <p>Use 0.01 level of significance to test whether there is a relationship between fidelity and selectivity. (Nov/Dec-2017)</p>	Fidelity				Selectivity	Low	Average	High	Low	6	12	32	Average	33	61	18	High	13	15	0	BTL -1	Remembering	PO1										
Fidelity																																		
Selectivity	Low	Average	High																															
Low	6	12	32																															
Average	33	61	18																															
High	13	15	0																															
2. (a)	A sample of 100 students is taken from a large population. The mean height of the students in this sample is 160cms. Can it be reasonably regarded that this sample is from a population of mean 165 cm and standard deviation 10 cm? Also estimate the 95% fiducial limits for the mean. (Apr/May-2015)	BTL -1	Remembering	PO1																														
2.(b)	<p>Given the following table for hair color and eye color, identify the value of Chi-square. Is there good association between hair color and eye color? (Nov/Dec-2012)</p> <table border="1"> <thead> <tr> <th colspan="2" rowspan="2"></th> <th colspan="3">Hair color</th> <th rowspan="2">Total</th> </tr> <tr> <th>Fair</th> <th>Brown</th> <th>Black</th> </tr> </thead> <tbody> <tr> <th rowspan="4">Eye color</th> <th>Blue</th> <td>15</td> <td>5</td> <td>20</td> <td>40</td> </tr> <tr> <th>Grey</th> <td>20</td> <td>10</td> <td>20</td> <td>50</td> </tr> <tr> <th>Brown</th> <td>25</td> <td>15</td> <td>20</td> <td>60</td> </tr> <tr> <th>Total</th> <td>60</td> <td>30</td> <td>60</td> <td>150</td> </tr> </tbody> </table>			Hair color			Total	Fair	Brown	Black	Eye color	Blue	15	5	20	40	Grey	20	10	20	50	Brown	25	15	20	60	Total	60	30	60	150	BTL -1	Remembering	PO1
				Hair color				Total																										
		Fair	Brown	Black																														
Eye color	Blue	15	5	20	40																													
	Grey	20	10	20	50																													
	Brown	25	15	20	60																													
	Total	60	30	60	150																													
3. (a)	<p>Two independent samples of sizes 8 and 7 contained the following values. (Nov/Dec-2013)</p> <table border="1"> <tbody> <tr> <td>Sample I</td> <td>19</td> <td>17</td> <td>15</td> <td>21</td> <td>16</td> <td>18</td> <td>16</td> <td>14</td> </tr> <tr> <td>Sample II</td> <td>15</td> <td>14</td> <td>15</td> <td>19</td> <td>15</td> <td>18</td> <td>16</td> <td></td> </tr> </tbody> </table> <p>(Apr/May-2016)</p>	Sample I	19	17	15	21	16	18	16	14	Sample II	15	14	15	19	15	18	16		BTL -2	Understanding	PO2												
Sample I	19	17	15	21	16	18	16	14																										
Sample II	15	14	15	19	15	18	16																											
3.(b)	<p>The following data gives the number of aircraft accidents that occurred during the various days of a week. Find whether the accidents are uniformly distributed over the week.</p> <table border="1"> <thead> <tr> <th>Days</th> <th>Su</th> <th>Mo</th> <th>Tue</th> <th>We</th> <th>Th</th> <th>Fr</th> <th>Sa</th> </tr> <tr> <th></th> <th>n</th> <th>n</th> <th>s</th> <th>d</th> <th>u</th> <th>i</th> <th>t</th> </tr> </thead> <tbody> <tr> <td>No. of accidents</td> <td>14</td> <td>16</td> <td>8</td> <td>12</td> <td>11</td> <td>9</td> <td>14</td> </tr> </tbody> </table>	Days	Su	Mo	Tue	We	Th	Fr	Sa		n	n	s	d	u	i	t	No. of accidents	14	16	8	12	11	9	14	BTL -2	Understanding	PO2						
Days	Su	Mo	Tue	We	Th	Fr	Sa																											
	n	n	s	d	u	i	t																											
No. of accidents	14	16	8	12	11	9	14																											

4. (a)	<p>Two independent samples of 8 and 7 items respectively had the following Values of the variable(weight in kgs) Use 0.05 LOS to test</p> <table border="1" data-bbox="250 205 881 289"> <tr> <td>Sample I</td> <td>9</td> <td>11</td> <td>13</td> <td>11</td> <td>15</td> <td>9</td> <td>12</td> <td>14</td> </tr> <tr> <td>Sample II</td> <td>10</td> <td>12</td> <td>10</td> <td>14</td> <td>9</td> <td>8</td> <td>10</td> <td></td> </tr> </table> <p>whether the variances of the two population's sample are equal. (Apr/May-2014)</p>	Sample I	9	11	13	11	15	9	12	14	Sample II	10	12	10	14	9	8	10		BTL -2	Understanding	PO2																		
Sample I	9	11	13	11	15	9	12	14																																
Sample II	10	12	10	14	9	8	10																																	
5. (a)	<p>A group of 10 rats fed on diet A and another group of 8 rats fed on diet B. Recorded the following increase the following increase in weight.(gm)</p> <table border="1" data-bbox="250 472 849 556"> <tr> <td>Diet A</td> <td>5</td> <td>6</td> <td>8</td> <td>1</td> <td>12</td> <td>4</td> <td>3</td> <td>9</td> <td>6</td> <td>10</td> </tr> <tr> <td>Diet B</td> <td>2</td> <td>3</td> <td>6</td> <td>8</td> <td>10</td> <td>1</td> <td>2</td> <td>8</td> <td>-</td> <td>-</td> </tr> </table> <p>Find the variances are significantly different. (Use F-test) (Nov/Dec-2014)</p>	Diet A	5	6	8	1	12	4	3	9	6	10	Diet B	2	3	6	8	10	1	2	8	-	-	BTL -2	Understanding	PO2														
Diet A	5	6	8	1	12	4	3	9	6	10																														
Diet B	2	3	6	8	10	1	2	8	-	-																														
5 (b)	<p>The marks obtained by a group of 9 regular course students and another group of 11 part time course students in a test are given below:</p> <table border="1" data-bbox="250 737 953 884"> <tr> <td>Sample I</td> <td>5</td> <td>6</td> <td>6</td> <td>5</td> <td>6</td> <td>5</td> <td>6</td> <td>6</td> <td>5</td> <td></td> <td></td> </tr> <tr> <td>Sample II</td> <td>6</td> <td>7</td> <td>7</td> <td>6</td> <td>6</td> <td>5</td> <td>7</td> <td>6</td> <td>7</td> <td>6</td> <td>6</td> </tr> <tr> <td></td> <td>2</td> <td>0</td> <td>1</td> <td>2</td> <td>0</td> <td>6</td> <td>5</td> <td>4</td> <td>2</td> <td>8</td> <td>6</td> </tr> </table> <p>Examine whether the marks obtained by regular students and part - time students differ significantly at 5% and 1% levels of significance. (Apr/May-2012)</p>	Sample I	5	6	6	5	6	5	6	6	5			Sample II	6	7	7	6	6	5	7	6	7	6	6		2	0	1	2	0	6	5	4	2	8	6	BTL -4	Applying	PO12
Sample I	5	6	6	5	6	5	6	6	5																															
Sample II	6	7	7	6	6	5	7	6	7	6	6																													
	2	0	1	2	0	6	5	4	2	8	6																													
6. (a)	<p>Two independent samples of sizes 8 and 7 contained the following values.</p> <table border="1" data-bbox="250 1066 909 1213"> <tr> <td>Sample I</td> <td>19</td> <td>17</td> <td>15</td> <td>21</td> <td>16</td> <td>18</td> <td>16</td> <td>14</td> </tr> <tr> <td>Sample II</td> <td>15</td> <td>14</td> <td>15</td> <td>19</td> <td>15</td> <td>18</td> <td>16</td> <td></td> </tr> </table> <p>Test if the two populations have the same variance.</p>	Sample I	19	17	15	21	16	18	16	14	Sample II	15	14	15	19	15	18	16		BTL -2	Understanding	PO2																		
Sample I	19	17	15	21	16	18	16	14																																
Sample II	15	14	15	19	15	18	16																																	
6.(b)	<p>In a certain factory there are two independent processes manufacturing the same item. The average weight in a sample of 250 items produced from one process is found to be 120 Ozs, with a standard deviation of 12 Ozs, while the corresponding figures in a sample of 400 items from the other process are 124 and 14. Is the difference between the two sample means significant? (Nov/Dec-2014)</p>	BTL -4	Applying	PO12																																				
7. (a)	<p>Records taken of the number of male and female births in 800 families having four Children are as follows :</p> <table data-bbox="250 1619 938 1724"> <tr> <td>Number of male births</td> <td>: 0</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> </tr> <tr> <td>Number of female births</td> <td>: 4</td> <td>3</td> <td>2</td> <td>1</td> <td>0</td> </tr> <tr> <td>Number of Families</td> <td>: 32</td> <td>178</td> <td>290</td> <td>236</td> <td>64</td> </tr> </table> <p>Infer whether the data are consistent with the hypothesis that the binomial law holds the chance of a male birth is equal to female birth, namely $p = \frac{1}{2} = q$. (Apr/May-2014)</p>	Number of male births	: 0	1	2	3	4	Number of female births	: 4	3	2	1	0	Number of Families	: 32	178	290	236	64	BTL -4	Analyzing	PO1,PO2, PO5																		
Number of male births	: 0	1	2	3	4																																			
Number of female births	: 4	3	2	1	0																																			
Number of Families	: 32	178	290	236	64																																			

7. (b)	<p>Samples of two types of electric bulbs were tested for length of life and following data were obtained.</p> <table border="1" data-bbox="370 170 881 317"> <tr> <td></td> <td>Type - I</td> <td>Type - II</td> </tr> <tr> <td>Sample size</td> <td>8</td> <td>7</td> </tr> <tr> <td>Sample mean</td> <td>1234 hrs</td> <td>1036 hrs</td> </tr> <tr> <td>Sample S.D</td> <td>36 hrs</td> <td>40 hrs</td> </tr> </table> <p>Analyze that, is the difference in the means sufficient to warrant that type I is superior to type II regarding the length of life? (Nov/Dec-2015)</p>		Type - I	Type - II	Sample size	8	7	Sample mean	1234 hrs	1036 hrs	Sample S.D	36 hrs	40 hrs	BTL -3	Applying	PO12									
	Type - I	Type - II																							
Sample size	8	7																							
Sample mean	1234 hrs	1036 hrs																							
Sample S.D	36 hrs	40 hrs																							
8. (a)	<p>A survey of 320 families with 5 children each revealed the following distribution</p> <table border="1" data-bbox="250 501 836 651"> <tr> <td>Boys</td> <td>5</td> <td>4</td> <td>3</td> <td>2</td> <td>1</td> <td>0</td> </tr> <tr> <td>Girls</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> </tr> <tr> <td>Families</td> <td>14</td> <td>56</td> <td>110</td> <td>88</td> <td>40</td> <td>12</td> </tr> </table> <p>Is this result consistent with the hypothesis that male and female births are equally probable?</p>	Boys	5	4	3	2	1	0	Girls	0	1	2	3	4	5	Families	14	56	110	88	40	12	BTL -6	Creating	PO1,PO2, PO12
Boys	5	4	3	2	1	0																			
Girls	0	1	2	3	4	5																			
Families	14	56	110	88	40	12																			
8.(b)	<p>The mean produce of wheat from a sample of 100 fields comes to 200kg per acre and another sample of 150 fields gives a mean 220 kg per acre. Assuming the standard deviation of the yield at 11 kg for the universe, test if there is a significant difference between the means of the samples? (Apr/May-2015)</p>	BTL -2	Understanding	PO2																					
9. (a)	<p>Two independent samples of sizes 9 and 7 from a normal population had the following values of the variables</p> <p>Sample 1 18 13 12 15 12 14 16 14 15</p> <p>Sample 2 16 19 13 16 18 13 15</p> <p>Justify whether the difference between the means of samples of samples significant? (Nov/Dec-2016)</p>	BTL -1	Remembering	PO1																					
9.(b)	<p>A simple sample of heights of 6400 Englishmen has a mean of 170cms and a standard deviation of 6.4cms, while a simple sample of heights of 1600 Americans has a mean of 172 cms and a standard deviation of 6.3cms. Do the data indicate that Americans are, on the average, taller than Englishmen?(BTL4) (Apr/May-2016)</p>	BTL -1	Remembering	PO1																					
10.(a)	<p>Two random samples gave the following results:</p> <table border="1" data-bbox="250 1486 904 1682"> <tr> <th>Sample</th> <th>Size</th> <th>Sample Mean</th> <th>Sum of squares of deviation from the mean</th> </tr> <tr> <td>1</td> <td>10</td> <td>15</td> <td>90</td> </tr> <tr> <td>2</td> <td>12</td> <td>14</td> <td>108</td> </tr> </table> <p>Analyze whether the samples have come from the same normal population. (Nov/Dec-2013)</p>	Sample	Size	Sample Mean	Sum of squares of deviation from the mean	1	10	15	90	2	12	14	108	BTL -1	Remembering	PO1									
Sample	Size	Sample Mean	Sum of squares of deviation from the mean																						
1	10	15	90																						
2	12	14	108																						
10.(b)	<p>A certain medicine administered to each of 10 patients resulted in the following increases in the B.P. 8, 8, 7, 5, 4, 1, 0, 0, -1, -1. Can it be concluded that the medicine was responsible for the increase in B.P. 5% I.o.s (Apr/May-2012)</p>	BTL -1	Remembering	PO1																					

11.(a)	200 digits were chosen at random from a set of tables. The frequencies of the digits were given below: Use χ^2 test to access the correctness of hypothesis that the digits were distributed in equal nos. in the table, given that the values of χ^2 are 16.9, 18.3, 19.7 for 9, 10, 11 degrees of freedom at 5% level of significance.	BTL -3	Applying	PO12																
11.(b)	4 coins were tossed 160 times and the following results were obtained No. of heads : 0 1 2 3 4 Observed frequencies: 17 52 54 31 6 Under the assumption that the coins are balanced, find the expected frequencies of getting 0, 1, 2, 3, 4 heads and test the goodness of fit. (Nov/Dec-2014)	BTL -5	Evaluating	PO1,PO2, PO5																
12.(a)	A sample of 200 persons with a particular disease was selected. Out of these, 100 were given a drug and the others were not given any drug. The result are as follows: <table border="1" style="margin-left: 20px;"> <thead> <tr> <th></th> <th>Drug s</th> <th>No Drugs</th> <th>Total</th> </tr> </thead> <tbody> <tr> <td>Cured</td> <td>65</td> <td>55</td> <td>120</td> </tr> <tr> <td>Not Cured</td> <td>35</td> <td>45</td> <td>80</td> </tr> <tr> <td>Total</td> <td>100</td> <td>100</td> <td>200</td> </tr> </tbody> </table> Test whether the drug is effective or not? (Apr/May-2015)		Drug s	No Drugs	Total	Cured	65	55	120	Not Cured	35	45	80	Total	100	100	200	BTL -1	Remembering	PO1
	Drug s	No Drugs	Total																	
Cured	65	55	120																	
Not Cured	35	45	80																	
Total	100	100	200																	
12.(b)	A certain stimulus administered to each of 12 patients resulted in the following increase of blood pressure 5,2,8,-1,3,0,-2,1,5,0, 4 & 6 can it be concluded that the stimulus will, in general, be accompanied by an increase in blood pressure?	BTL -6	Creating	PO1,PO2, PO12																
13.(a)	In a referendum submitted by the students to the body at a university, 850 men and 560 women voted. 500 men and 320 women voted favorably. Does this indicate a significant difference of opinion between men and women on this matter at 1% level of significance? (Nov/Dec-2014)	BTL -1	Remembering	PO1																
13.(b)	Random samples drawn from two places gave the following data relating to the heights of male adults: <table border="1" style="margin-left: 20px;"> <thead> <tr> <th></th> <th>Place A</th> <th>Place B</th> </tr> </thead> <tbody> <tr> <td>Mean height</td> <td>8</td> <td>7</td> </tr> <tr> <td>S.D</td> <td>1234 hrs</td> <td>1036 hrs</td> </tr> <tr> <td>No. of sample</td> <td>36 hrs</td> <td>40 hrs</td> </tr> </tbody> </table> Test at 5 % level, that the mean height is the same for adults in the two places. (Apr/May-2012)		Place A	Place B	Mean height	8	7	S.D	1234 hrs	1036 hrs	No. of sample	36 hrs	40 hrs	BTL -2	Understanding	PO2				
	Place A	Place B																		
Mean height	8	7																		
S.D	1234 hrs	1036 hrs																		
No. of sample	36 hrs	40 hrs																		
14.(a)	In a random sample of 1000 people from city A, 400 are found to be consumers of rice. In a sample of 800 from city B, 400 are found to be consumers of rice. Does this data give a significant difference between the two cities as far as the proportion of rice consumers is concerned?	BTL -4	Analyzing	PO1,PO2, PO5																

	(Nov/Dec-2012)			
14.(b)	In a year there are 956 births in a town A of which 52.5% were male while in towns A and B combined, this proportion in a total of 1406 births was 0.496. Is there any significant difference in the proportion of male births in the two towns ? (Apr/May-2011)	BTL -2	Understanding	PO2

UNIT - II DESIGN OF EXPERIMENTS				
One way and two way classifications - Completely randomized design - Randomized block design - Latin square design - 2 ² factorial design.				
PART - A				
Q.No	Question	BT Level	Competence	PO
1.	Write the advantages of Latin Square (Nov/Dec-2017)	BTL -1	Remembering	PO1
2.	What are the conditions to be followed in one way classification?(Nov/Dec-2017)	BTL -1	Remembering	PO1
3.	What is meant by analysis of variance?(May/Jun-2016)	BTL -1	Remembering	PO1
4.	Why a 2x2 Latin square is not possible? Explain.(May/Jun-2016)(May/Jun-2014).	BTL -1	Remembering	PO1
5.	Define Replication and Randomization.(Nov/Dec-	BTL -1	Remembering	PO1
6.	What is the advantage of factorial experiment? (Nov/Dec-2016)	BTL -1	Remembering	PO1
7.	What is the aim of design of experiment?(Apr/May-2015)	BTL -2	Understanding	PO2
8.	What are the basic principles of experimental design? (Apr/May-2015)	BTL -2	Understanding	PO2
9.	Write the advantages and disadvantages of RBD?(Apr/May-2015)	BTL -2	Understanding	PO2
10.	What is Latin Square design ?	BTL -2	Understanding	PO2
11.	Define Raw Sum of Squares and Correction factor	BTL -3	Applying	PO1,PO2,PO12
12.	Write any 3 applications of LSD. (Nov/Dec-2014)	BTL -3	Applying	PO1,PO2,PO12
13.	How do you calculate the Correction factor in LSD? (Nov/Dec-2012)	BTL -3	Applying	PO1,PO2,PO12
14.	What do you mean by design of experiments?(Nov/Dec-2014)	BTL -4	Analyzing	PO5

15.	What are the subject matters included in the design of experiment?	BTL -4	Analyzing	PO5
16.	What are the assumptions in ANOVA? (Apr/May-	BTL -4	Analyzing	PO5
17.	are the three essential steps to plan an experiment?	BTL -5	Evaluating	PO1,PO2,PO5
18.	What are the basic steps in ANOVA? (Apr/May-2014)	BTL -5	Evaluating	PO1,PO2,PO5
19.	Write the steps to find F-ratio. (Nov/Dec-2016)	BTL -6	Creating	PO1,PO2,PO5
20.	Discuss the advantages of Completely Randomized block design.	BTL -6	Creating	PO1,PO2,PO5
21.	State the uses of ANOVA. (Apr/May-2015)	BTL -4	Analyzing	PO12
22.	Explain the word treatment in ANOVA. (Apr/May-2015)	BTL -4	Analyzing	PO12
23.	What do you mean by 2-way classification?	BTL -4	Analyzing	PO12
24.	Indicate the characteristics of a good experimental Design (Nov/Dec-2011)	BTL -5	Evaluating	PO1,PO2,PO5
25.	What are the important designs of experiments?	BTL -5	Evaluating	PO1,PO2,PO5
26.	What is an experimental error? (Nov/Dec-2011)	BTL -6	Creating	PO1,PO2,PO5
27.	What is meant by CRD? (Apr/May-2012)	BTL -6	Creating	PO1,PO2,PO5
28.	Compare RBD and LSD.	BTL -3	Applying	PO1,PO2,PO5
29.	Compare LSD and RBD. (Apr/May-2015)	BTL -3	Applying	PO1,PO2,PO5
30.	What are the uses of Chi-Square test?	BTL -4	Analyzing	PO5
PART - B				
1.(a)	The accompanying data resulted from an experiment comparing the degree of soiling for fabric copolymerized with the 3 different mixtures of met acrylic acid. Analyse the classification. Mixture 1 : 0.56 1.12 0.90 1.07 0.94 Mixture 2 : 0.72 0.69 0.87 0.78 0.91 Mixture 3 : 0.62 1.08 1.07 0.99 0.93 (Apr/May-2017)	BTL -1	Remembering	PO1
1. (b)	A set of data involving 4 tropical food stuffs A, B, C, D tried on 20 chicks is given below. All the 20 chicks are treated alike in all respects except the feeding treatments and each feeding treatment is given to 5 chicks. Analyze the data: A 55 49 42 21 52 B 61 112 30 89 63 C 42 97 81 95 92 D 169 137 169 85 154 (Apr/May-2016)	BTL -2	Understanding	PO2

2. (a)	<p>The following table shows the lives in hours of four brands of electric lamps brand A:1610, 1610, 1650, 1680, 1700, 1720, 1800 B: 1580, 1640, 1640, 1700, 1750 C:1460, 1550, 1600, 1620, 1640, 1660, 1740, 1820 D:1510, 1520, 1530, 1570, 1600, 1680 Identify an analysis of variance and test the homogeneity of the mean lives of the four brands of lamps. (Apr/May-2014)</p>	BTL -1	Remembering	PO1																																										
2.(b)	<p>A company appoints 4 salesmen A, B, C and D and observes their sales in 3 seasons, summer winter and monsoon. The figures are given in the following table:</p> <table border="1" data-bbox="263 546 850 735"> <thead> <tr> <th></th> <th colspan="4">SALESMEN</th> </tr> <tr> <th>SEASON</th> <th>1</th> <th>2</th> <th>3</th> <th>4</th> </tr> </thead> <tbody> <tr> <td>SUMMER</td> <td>45</td> <td>40</td> <td>28</td> <td>37</td> </tr> <tr> <td>WINTER</td> <td>43</td> <td>41</td> <td>45</td> <td>38</td> </tr> <tr> <td>MONSOON</td> <td>39</td> <td>39</td> <td>43</td> <td>41</td> </tr> </tbody> </table> <p>Carry out an Analysis of variances. (Apr/May-2013)</p>		SALESMEN				SEASON	1	2	3	4	SUMMER	45	40	28	37	WINTER	43	41	45	38	MONSOON	39	39	43	41	BTL -2	Understanding	PO2																	
	SALESMEN																																													
SEASON	1	2	3	4																																										
SUMMER	45	40	28	37																																										
WINTER	43	41	45	38																																										
MONSOON	39	39	43	41																																										
3.	<p>In order to determine whether there is significant difference in the durability of 3 makes of computers, samples of size 5 are selected from each make and the frequency of repair during the first year of purchase is observed. The results are as follows: In view of the above data, what conclusion can you draw?</p> <table border="1" data-bbox="457 991 659 1255"> <thead> <tr> <th colspan="3">MAKES</th> </tr> <tr> <th>A</th> <th>B</th> <th>C</th> </tr> </thead> <tbody> <tr> <td>5</td> <td>8</td> <td>7</td> </tr> <tr> <td>6</td> <td>10</td> <td>3</td> </tr> <tr> <td>8</td> <td>11</td> <td>5</td> </tr> <tr> <td>9</td> <td>12</td> <td>4</td> </tr> <tr> <td>7</td> <td>4</td> <td>1</td> </tr> </tbody> </table> <p>(Apr/May-2012)</p>	MAKES			A	B	C	5	8	7	6	10	3	8	11	5	9	12	4	7	4	1	BTL -1	Remembering	PO1																					
MAKES																																														
A	B	C																																												
5	8	7																																												
6	10	3																																												
8	11	5																																												
9	12	4																																												
7	4	1																																												
4.	<p>Five doctors each test five treatments for a certain disease and observe the number of days each patient take store cover. The results are as follows (recovery time in days)</p> <table border="1" data-bbox="263 1438 850 1703"> <thead> <tr> <th></th> <th colspan="5">TREATMENT</th> </tr> <tr> <th>DOCTOR</th> <th>A</th> <th>B</th> <th>C</th> <th>D</th> <th>E</th> </tr> </thead> <tbody> <tr> <td>A</td> <td>10</td> <td>14</td> <td>23</td> <td>18</td> <td>20</td> </tr> <tr> <td>B</td> <td>11</td> <td>15</td> <td>24</td> <td>17</td> <td>21</td> </tr> <tr> <td>C</td> <td>9</td> <td>12</td> <td>20</td> <td>16</td> <td>19</td> </tr> <tr> <td>D</td> <td>8</td> <td>13</td> <td>17</td> <td>17</td> <td>20</td> </tr> <tr> <td>E</td> <td>12</td> <td>15</td> <td>19</td> <td>15</td> <td>22</td> </tr> </tbody> </table> <p>Estimate the difference between (a) doctors and (b) treatments for the above data at 5% level. (Nov/Dec-2017)</p>		TREATMENT					DOCTOR	A	B	C	D	E	A	10	14	23	18	20	B	11	15	24	17	21	C	9	12	20	16	19	D	8	13	17	17	20	E	12	15	19	15	22	BTL -2	Understanding	PO2
	TREATMENT																																													
DOCTOR	A	B	C	D	E																																									
A	10	14	23	18	20																																									
B	11	15	24	17	21																																									
C	9	12	20	16	19																																									
D	8	13	17	17	20																																									
E	12	15	19	15	22																																									
5.	<p>Perform a 2-way ANOVA on the data given below:</p> <table border="1" data-bbox="349 1848 764 1959"> <thead> <tr> <th colspan="2"></th> <th colspan="3">Treatment 1</th> </tr> <tr> <th colspan="2"></th> <th>1</th> <th>2</th> <th>3</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>1</td> <td>30</td> <td>26</td> <td>38</td> </tr> </tbody> </table>			Treatment 1					1	2	3	1	1	30	26	38	BTL -3	Applying	PO1,PO2,PO5																											
		Treatment 1																																												
		1	2	3																																										
1	1	30	26	38																																										

	<table border="1"> <tr><td>2</td><td>24</td><td>29</td><td>28</td></tr> <tr><td>3</td><td>33</td><td>24</td><td>35</td></tr> <tr><td>4</td><td>36</td><td>31</td><td>30</td></tr> <tr><td>5</td><td>27</td><td>35</td><td>33</td></tr> </table> <p>Use the coding method subtracting 30 from the given no. (Nov/Dec-2016)</p>	2	24	29	28	3	33	24	35	4	36	31	30	5	27	35	33																								
2	24	29	28																																						
3	33	24	35																																						
4	36	31	30																																						
5	27	35	33																																						
6.	<p>The following data represent a certain person to work from Monday to Friday by 4 different routes.</p> <table border="1"> <thead> <tr><th colspan="2" rowspan="2"></th><th colspan="5">DAYS</th></tr> <tr><th>MON</th><th>TUE</th><th>WED</th><th>THU</th><th>FRI</th></tr> </thead> <tbody> <tr><th rowspan="4">ROUTES</th><th>1</th><td>22</td><td>26</td><td>25</td><td>25</td><td>31</td></tr> <tr><th>2</th><td>25</td><td>27</td><td>28</td><td>26</td><td>29</td></tr> <tr><th>3</th><td>26</td><td>29</td><td>33</td><td>30</td><td>33</td></tr> <tr><th>4</th><td>26</td><td>28</td><td>27</td><td>30</td><td>30</td></tr> </tbody> </table> <p>Test at 5% level of significance whether the differences among the means obtained for the different routes are significant and also whether the differences among the means obtained for the different days of the week are significant. (Nov/Dec-2015)</p>			DAYS					MON	TUE	WED	THU	FRI	ROUTES	1	22	26	25	25	31	2	25	27	28	26	29	3	26	29	33	30	33	4	26	28	27	30	30	BTL -2	Understanding	PO2
				DAYS																																					
		MON	TUE	WED	THU	FRI																																			
ROUTES	1	22	26	25	25	31																																			
	2	25	27	28	26	29																																			
	3	26	29	33	30	33																																			
	4	26	28	27	30	30																																			
7.	<p>Analyze the variance in the following Latin square of yields of paddy where A, B, C, D denote the different methods of cultivation..</p> <table border="1"> <tr><td>D122</td><td>A121</td><td>C123</td><td>B122</td></tr> <tr><td>B124</td><td>C123</td><td>A122</td><td>D125</td></tr> <tr><td>A120</td><td>B119</td><td>D120</td><td>C121</td></tr> <tr><td>C122</td><td>D123</td><td>B121</td><td>A122</td></tr> </table> <p>Examine whether the different methods of cultivation have given significantly different yields. (Nov/Dec-2014)</p>	D122	A121	C123	B122	B124	C123	A122	D125	A120	B119	D120	C121	C122	D123	B121	A122	BTL -4	Analyzing	PO5																					
D122	A121	C123	B122																																						
B124	C123	A122	D125																																						
A120	B119	D120	C121																																						
C122	D123	B121	A122																																						
8. (a)	<p>The following data resulted from an experiment to compare three burners A, B, C. A Latin square design was used as the tests were made on 3 engines and were spread over 3 days.</p> <table border="1"> <tr><td>A16</td><td>B17</td><td>C20</td></tr> <tr><td>B16</td><td>C21</td><td>A15</td></tr> <tr><td>C15</td><td>A12</td><td>B13</td></tr> </table> <p>(Nov/Dec-2013)</p>	A16	B17	C20	B16	C21	A15	C15	A12	B13	BTL -1	Remembering	PO1																												
A16	B17	C20																																							
B16	C21	A15																																							
C15	A12	B13																																							
8.(b)	<p>A variable trial was conducted on wheat with 4 varieties in a Latin square design. The plan of the experiment and the per plot yield are given below.</p> <p>C25 B23 A20 D20 A19 D19 C21 B18 B19 A14 D17 C20 D17 C20 B21 A15</p> <p>(Nov/Dec-2012)</p>	BTL -5	Evaluating	PO5																																					

<p>9.</p>	<p>A farmer wishes to test the effects of four different fertilizers A, B, C, D on the yield of Wheat. In order to eliminate sources of error due to variability in soil fertility, he uses the fertilizers, in a Latin square arrangement as indicated in the following table, where the numbers indicate yields per unit area.</p> <table border="1" data-bbox="196 310 570 478"> <tr><td>A18</td><td>C21</td><td>D25</td><td>B11</td></tr> <tr><td>D22</td><td>B12</td><td>A15</td><td>C19</td></tr> <tr><td>B15</td><td>A20</td><td>C23</td><td>D24</td></tr> <tr><td>C22</td><td>D21</td><td>B10</td><td>A17</td></tr> </table> <p>Design an analysis of variance to determine if there is a significant difference between the fertilizers at $\alpha=0.05$ and $\alpha=0.01$ levels of significance.</p>	A18	C21	D25	B11	D22	B12	A15	C19	B15	A20	C23	D24	C22	D21	B10	A17	<p>BTL -1</p>	<p>Remembering</p>	<p>PO1</p>														
A18	C21	D25	B11																															
D22	B12	A15	C19																															
B15	A20	C23	D24																															
C22	D21	B10	A17																															
<p>10.</p>	<p>Set up the analysis of variance for the following results of a Latin Square Design(use $\alpha = 0.01$) level of significance</p> <table border="1" data-bbox="370 695 743 863"> <tr><td>A12</td><td>C19</td><td>B10</td><td>D8</td></tr> <tr><td>C18</td><td>B12</td><td>D6</td><td>A7</td></tr> <tr><td>B22</td><td>D10</td><td>A5</td><td>C21</td></tr> <tr><td>D12</td><td>A7</td><td>C27</td><td>B17</td></tr> </table>	A12	C19	B10	D8	C18	B12	D6	A7	B22	D10	A5	C21	D12	A7	C27	B17	<p>BTL -4</p>	<p>Analyzing</p>	<p>PO5</p>														
A12	C19	B10	D8																															
C18	B12	D6	A7																															
B22	D10	A5	C21																															
D12	A7	C27	B17																															
<p>11.</p>	<p>In a 5x5 Latin square experiment, the data collected is given in the matrix below Yield per plot is given in quintals for the five different cultivation treatments A, B, C,D and E. Perform the analysis of variance.</p> <p>A48 E66 D56 C52 B61 D64 B62 A50 E64 C63 B69 A53 C60 D61 E67 C57 D58 E67 B65 A55 E67 C57 B66 A60 D57</p>	<p>BTL -6</p>	<p>Creating</p>	<p>PO1,PO2,PO5</p>																														
<p>12</p>	<p>In a Latin square experiment given below are the yields in quintals per acre on the paddy crop carried out for testing the effect of five fertilizers A, B, C, D, E. Analyze the data for variations.</p> <p>B25 A18 E27 D30 C27 A19 D31 C29 E26 B23 C28 B22 D33 A18 E27 E28 C26 A20 B25 D33 D32 E25 B23 C28 A20</p>	<p>BTL -3</p>	<p>Applying</p>	<p>PO1,PO2,PO5</p>																														
<p>13.</p>	<p>Find out the main effects and interaction effects in the following 2² factorial experiment and write down the analysis of variance table</p> <table border="1" data-bbox="326 1682 792 1906"> <tr><td>BLOCKS</td><td>1</td><td>A</td><td>B</td><td>AB</td></tr> <tr><td></td><td>0</td><td>10</td><td>1</td><td>11</td></tr> <tr><td>I</td><td>64</td><td>25</td><td>30</td><td>60</td></tr> <tr><td>II</td><td>75</td><td>14</td><td>50</td><td>33</td></tr> <tr><td>III</td><td>76</td><td>12</td><td>41</td><td>17</td></tr> <tr><td>IV</td><td>75</td><td>33</td><td>25</td><td>10</td></tr> </table>	BLOCKS	1	A	B	AB		0	10	1	11	I	64	25	30	60	II	75	14	50	33	III	76	12	41	17	IV	75	33	25	10	<p>BTL -3</p>	<p>Applying</p>	<p>PO1,PO2,PO5</p>
BLOCKS	1	A	B	AB																														
	0	10	1	11																														
I	64	25	30	60																														
II	75	14	50	33																														
III	76	12	41	17																														
IV	75	33	25	10																														

14.	<p>An experiment was planned to study the effect of sulphate of potash and super phosphate on the yields of potatoes. All the combinations of 2 levels of super phosphate (p) and two levels of sulphate (k) of potash were studied in a RBD with 4 replication for each. The yields obtained are given in the following table. The yields obtained are given in the following table. Analyze the data and give your conclusion (with $\alpha = 1\%$)</p>																																												
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 15%;">BLOCKS</th> <th colspan="4">Yields (Per Plot)</th> </tr> </thead> <tbody> <tr> <td rowspan="2" style="text-align: center;">I</td> <td style="text-align: center;">(1)</td> <td style="text-align: center;">a</td> <td style="text-align: center;">b</td> <td style="text-align: center;">ab</td> </tr> <tr> <td style="text-align: center;">23</td> <td style="text-align: center;">25</td> <td style="text-align: center;">22</td> <td style="text-align: center;">38</td> </tr> <tr> <td rowspan="2" style="text-align: center;">II</td> <td style="text-align: center;">P</td> <td style="text-align: center;">(1)</td> <td style="text-align: center;">K</td> <td style="text-align: center;">KP</td> </tr> <tr> <td style="text-align: center;">40</td> <td style="text-align: center;">26</td> <td style="text-align: center;">36</td> <td style="text-align: center;">38</td> </tr> <tr> <td rowspan="2" style="text-align: center;">III</td> <td style="text-align: center;">(1)</td> <td style="text-align: center;">K</td> <td style="text-align: center;">KP</td> <td style="text-align: center;">P</td> </tr> <tr> <td style="text-align: center;">29</td> <td style="text-align: center;">20</td> <td style="text-align: center;">30</td> <td style="text-align: center;">20</td> </tr> <tr> <td rowspan="2" style="text-align: center;">IV</td> <td style="text-align: center;">KP</td> <td style="text-align: center;">K</td> <td style="text-align: center;">P</td> <td style="text-align: center;">(1)</td> </tr> <tr> <td style="text-align: center;">34</td> <td style="text-align: center;">31</td> <td style="text-align: center;">24</td> <td style="text-align: center;">28</td> </tr> </tbody> </table>	BLOCKS	Yields (Per Plot)				I	(1)	a	b	ab	23	25	22	38	II	P	(1)	K	KP	40	26	36	38	III	(1)	K	KP	P	29	20	30	20	IV	KP	K	P	(1)	34	31	24	28	BTL -3	Applying	PO1,PO2,PO1 2
BLOCKS	Yields (Per Plot)																																												
I	(1)	a	b	ab																																									
	23	25	22	38																																									
II	P	(1)	K	KP																																									
	40	26	36	38																																									
III	(1)	K	KP	P																																									
	29	20	30	20																																									
IV	KP	K	P	(1)																																									
	34	31	24	28																																									

UNIT III SOLUTION OF EQUATIONS AND EIGENVALUE PROBLEMS

Newton Raphson method - Gauss elimination method - pivoting - Gauss Jordan methods - Iterative methods of Gauss Jacobi and Gauss Seidel - Matrix inversion by Gauss Jordan method - Eigen values of a matrix by power method.
 Textbook : Grewal. B.S., and Grewal. J.S., "Numerical Methods in Engineering and Science", 9th Edition, Khanna Publishers, New Delhi, 2007.

PART - A

Q.No	Questions	BT Level	Competence	PO
1	State the order (rate) of convergence and convergence condition for Newton Raphson method. (A.U.N/D 2017, N/D 2011,2012, M/J 2013)	BTL-4	Analyzing	PO1
2	Give Newton Raphson iterative formula. (A.U N/D 2009,M/J 2012,2014)	BTL-2	Understanding	PO1,PO2 ,PO3
3	Establish an iteration formula to find the reciprocal of a positive number N by Newton Raphson method. (A.U.N/D 2010, M/J 2012)	BTL-1	Remembering	PO1,PO2
4	State the principle used in Gauss-Jordan method. (A.U M/J 2011)	BTL-1	Remembering	PO1
5	Give the sufficient condition of convergence of Gauss Seidel method. . (A.U M/J 2011)	BTL-1	Remembering	PO1
6	Write the conditions for convergence in Gauss Seidel iterative technique. (or) When the method of iteration will be useful ? (A.U M/J 2009)	BTL-3	Applying	PO1
7	State Gauss Seidel method. (A.U M/J 2011,N/D 2012)	BTL-1	Remembering	PO1,PO2
8	Gauss Seidel method always converges - True or False. . (A.U	BTL-1	Remembering	PO1,PO2

	M/J 2016)			
9	Write the first iteration values of x,y,z when the equations $27x+6y-z = 85$, $6x+15y+2z = 72$, $x+y+5z = 110$ are solved by Gauss Seidel method. (A.U N/D 2009,M/J 2012,2016)	BTL-3	Applying	PO1
10	Compare Gauss Elimination and Gauss Jordan methods for solving linear systems of the form $AX=B$. (A.U M/J 2016)	BTL-1	Remembering	PO1
11	What type of Eigen value can be obtained using power method? (A.U.N/D 2017, N/D 2011,2012, M/J 2014)	BTL-1	Remembering	PO1
12	Find the dominant eigen value of $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ by power method. (A.U M/J 2012)	BTL-1	Remembering	PO1
13	On what type of equations Newton's method can be applicable ? (A.U A/M 2016)	BTL-1	Remembering	PO1,PO2 ,PO5
14	By Gauss elimination method solve $x + y = 2$ and $2x + 3y = 5$. (A.U M/J 2014)	BTL-1	Remembering	PO1
15	Why Gauss Seidel iteration is a method of successive corrections? (A.U M/J 2016)	BTL-4	Analyzing	PO1
16	What are the merits of Newton's method of iteration?	BTL-1	Remembering	PO1
17	Give two direct methods to solve a system of linear equations. . (A.U A/M 2013)	BTL-2	Understanding	PO2
18	Compare Gauss Elimination with Gauss Seidel method.(A.U M/J 2017)	BTL-1	Remembering	PO1
19	Find inverse of $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ by Gauss Jordan method. (A.U M/J 2013)	BTL-1	Remembering	PO1,PO2

PART-B

1	Solve $x \log_{10} x = 12.34$ with $x_0 = 10$ using Newton's method. (A.U.N/D 2017, N/D 2011,2012, M/J 2013)	BTL-4	Analyzing	PO1,PO2 ,PO5
2	Find the negative root of the equation $\sin x = 1 + x^3$ by using Newton Raphson method. (A.U M/J 2015)	BTL-4	Analyzing	PO1,PO2 ,PO5
3	Solve the following equation by Gauss Elimination method $10x - 2y + 3z = 23$ $2x + 10y - 5z = -33$ (A.U.N/D 2017, N/D 2011,2012, M/J 2014) $3x - 4y + 10z = 41$	BTL-5	Evaluating	PO1,PO2 ,PO5
4	Solve the equation by Gauss Jordan method : $2x_1 + x_2 + 4x_3 = 4$ $x_1 - 3x_2 - x_3 = -5$ $3x_1 - 2x_2 + 2x_3 = -1$	BTL-5	Evaluating	PO1,PO2 ,PO5

5	Find the inverse of $\begin{bmatrix} 2 & 2 & 3 \\ 2 & 1 & 1 \\ 1 & 3 & 5 \end{bmatrix}$ using Gauss Jordan method.	BTL-2	Understanding	PO1,PO2
6	Solve by Gauss Siedel method $x + y + 54z = 110$ $27x + 6y - z = 85$ $6x + 15y + 2z = 72$ (A.U.N/D 2017, N/D 2011,2013, M/J 2014)	BTL-2	Understanding	PO1,PO2
7	Find the dominant (largest) eigen value and the corresponding eigen vector of $A = \begin{bmatrix} 1 & -3 & 2 \\ 4 & 4 & -1 \\ 6 & 3 & 5 \end{bmatrix}$ by power method. (A.U M/J 2015)	BTL-5	Evaluating	PO1,PO2 ,PO5
8	Find the numerically largest eigen value of $A = \begin{bmatrix} 25 & 1 & 2 \\ 1 & 3 & 0 \\ 2 & 0 & -4 \end{bmatrix}$ by power method and the corresponding eigen vector. (A.U M/J 2011,N/D 2012)	BTL-5	Evaluating	PO1,PO2 ,PO5
9	Find the numerically largest eigen value of $A = \begin{bmatrix} 5 & 4 & 3 \\ 10 & 8 & 6 \\ 20 & -4 & 22 \end{bmatrix}$ by power method with the initial eigen vector $X_0 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$. (A.U M/J 2016)	BTL-5	Evaluating	PO1,PO2 ,PO5

UNIT IV INTERPOLATION, NUMERICAL DIFFERENTIATION AND NUMERICAL INTEGRATION

Lagrange's and Newton's divided difference interpolations - Newton's forward and backward difference interpolation - Approximation of derivates using interpolation polynomials - Numerical

single and double integrations using Trapezoidal and Simpson's 1/3 rules.
 Textbook : Grewal. B.S., and Grewal. J.S., "Numerical Methods in Engineering and Science", 9th Edition, Khanna Publishers, New Delhi, 2007.

PART - A

CO Mapping : C214.2

Q. No	Questions	BT Level	Competence	PO
1	Define interpolation and extrapolation? (A.U.N/D 2017, N/D 2011,2012, M/J 2013)	BTL-4	Analyzing	PO1
2	State Newton's formula on interpolation. When it is used? (A.U.N/D 2017, N/D 2011,2012, M/J 2014)	BTL-1	Remembering	PO1,PO2
3	Say True or False. - Newton's divided difference formula is applicable only for equally spaced intervals. (A.U M/J 2011)	BTL-2	Understanding	PO1,PO2
4	State Newton's divided difference formula.	BTL-4	Analyzing	PO2
5	State Lagrange's interpolation formula	BTL-1	Remembering	PO1
6	Use Lagrange's formula to find the quadratic polynomial that takes these values $\begin{matrix} x : & 0 & 1 & 3 \\ y : & 0 & 1 & 0 \end{matrix}$ Then find y(2). (A.U M/J 2011,N/D 2012)	BTL-2	Understanding	PO1
7	By differentiating Newton forward and backward difference formula, find the first derivative of the function f(x). (A.U M/J 2013)	BTL-2	Understanding	PO1,PO2
8	Write down the Newton - cotes quadrature formula.	BTL-1	Remembering	PO1
9	What is the geometrical interpretation of Trapezoidal rule? (A.U M/J 2016,N/D 2012)	BTL-1	Remembering	PO1
10	Using Trapezoidal rule evaluate $\int_0^{\pi} \sin x dx$ by dividing the range into 6 equal parts.	BTL-1	Remembering	PO1
11	Why is Trapezoidal rule so called? (A.U N/D 2011,N/D 2014)	BTL-2	Understanding	PO1,PO2
12	What are the truncation errors in Trapezoidal and Simpson's rules of numerical integration?	BTL-4	Analyzing	PO1
13	What is the condition for Simpson's 3/8 rule and state the formula.	BTL-4	Analyzing	PO1,PO2
14	Using Simpson's rule find $\int_0^4 e^x dx$ given $e^0 = 1, e^1 = 2.72, e^2 = 7.39, e^3 = 20.09, e^4 = 54.6$	BTL-4	Analyzing	PO1
15	Compare Trapezoidal rule and Simpson's 1/3 rd rule for evaluating numerical integration. (A.U M/J 2015,N/D 2017)	BTL-1	Remembering	PO1

PART - B

1	Construct Newton's forward interpolation polynomial for the following data. $x: 4 \quad 6 \quad 8 \quad 10$ $y: 1 \quad 3 \quad 8 \quad 16$ Use it to find the value of y for $x = 5$. (A.U M/J 2011,A/M 2012)	BTL-5	Evaluating	PO1,PO2 PO3,PO5
2	The following data are taken from the steam table $\text{Temp}^{\circ} \text{c}: 140 \quad 150 \quad 160 \quad 170 \quad 180$ Pressure kg $\text{f/cm}^2: 3.685 \quad 4.854 \quad 6.302 \quad 8.076 \quad 10.225$ Find the pressure at temperature $t = 175^{\circ}$.	BTL-4	Analyzing	PO1,PO2
3	Using Lagrange's interpolation formula calculate the profit in the year 2000 from the following data $\text{Year}: 1997 \quad 1999 \quad 2001 \quad 2002$ $\left. \begin{array}{l} \text{Profit in lakhs} \\ \text{of Rs.} \end{array} \right\} : 43 \quad 65 \quad 159 \quad 248$ (A.U.N/D 2017, N/D 2011,2012, M/J 2013)	BTL-5	Evaluating	PO1,PO2, PO5,PO12
4	Find the polynomial $f(x)$ by using Lagrange's formula and hence find $f(3)$ for $x: 0 \quad 1 \quad 2 \quad 5$ $f(x): 2 \quad 3 \quad 12 \quad 147$ (A.U.N/D 2017, N/D 2011,2014, M/J 2013)	BTL-4	Analyzing	PO1,PO2, PO5,PO12
5	Using Newton divided difference formula find $u(3)$ given $u(1) = -26, u(2) = 12, u(4) = 256, u(6) = 844$.	BTL-5	Evaluating	PO1,PO2, PO5,PO12
6	From the given table, the values of y are consecutive terms of a series of which 23.6 is the sixth term. Find the first and tenth terms of the series. $x: 3 \quad 4 \quad 5 \quad 6 \quad 7 \quad 8 \quad 9$ $y: 4.8 \quad 8.4 \quad 14.5 \quad 23.6 \quad 36.2 \quad 52.8 \quad 73.9$ (A.U M/J 2016)	BTL-4	Analyzing	PO1,PO2, PO5,PO12
7	The following data gives the velocity of a particle for 20 seconds at an interval of 5 seconds. Find the initial acceleration using the entire data $\text{time (sec.)}: 0 \quad 5 \quad 10 \quad 15 \quad 20$ $\text{velocity (m/sec.)}: 0 \quad 3 \quad 14 \quad 69 \quad 228$ (A.U N/D 2015)	BTL-5	Evaluating	PO1,PO2, PO5,PO12
8	Using Trapezoidal rule, evaluate $\int_{-1}^1 \frac{dx}{1+x^2}$ taking 8 intervals.	BTL-5	Evaluating	PO1,PO2, PO5,PO12

9	Find an approximate value of $\log_e 5$ by calculating to four decimal places by Simpson's rule the integral $\int_0^5 \frac{dx}{4x+5}$ dividing the range into 10 equal parts. (A.U A/M 2016)	BTL-3	Applying	PO1,PO2, PO5,PO12
10	Evaluate $\int_0^6 \frac{dx}{1+x^2}$ by dividing the range into 6 equal parts using Simpson's rule.	BTL-3	Applying	PO1,PO2, PO5,PO12
11.	Evaluate $\int_0^1 \frac{dx}{1+x^2}$ take $h = 0.125$. Hence find π using Simpson's rule. (A.U.N/D 2017, N/D 2011,2012, M/J 2014)	BTL-5	Evaluating	PO1,PO2, PO5,PO12
12.	Compute $\int_0^1 \frac{xdx}{x^3+10}$ using Trapezoidal rule and Simpson's rule with the number of points 3,5,9. (A.U M/J 2017)	BTL-3	Applying	PO1,PO2, PO5,PO12

UNIT V NUMERICAL SOLUTION OF ORDINARY DIFFERENTIAL EQUATIONS

Taylor's series method - Euler's method - Modified Euler's method - Fourth order Runge-Kutta method for solving first and second order equations - Milne's predictor-corrector methods for solving first order equations - Finite difference methods for solving second order equation.

PART – A

CO Mapping :

Q.No	Questions	BT Level	Competence	PO
1.	State Modified Euler algorithm to solve $y' = f(x, y), y(x_0) = y_0$ at $x=x_0+h$. (A.U.N/D 2017, N/D 2011,2012, M/J 2013)	BTL -1	Remembering	PO1
2.	State the disadvantage of Taylor series method. (A.U N/D 2009,M/J 2012,2014)	BTL -1	Understanding	PO1
3.	Write the merits and demerits of the Taylor method of solution. (A.U.N/D 2010, M/J 2012)	BTL -5	Understanding	PO1
4.	Which is better Taylor's method or R. K. Method?(or)	BTL -1	Remembering	PO1

	State the special advantage of Runge-Kutta method over Taylor series method. (A.U M/J 2011)			
5.	Compare Runge-Kutta methods and predictor – corrector methods for solution of initial value problem. (A.U M/J 2011)	BTL -1	Remembering	PO1
6.	What is a Predictor-corrector method of solving a differential equation?(A.U M/J 2009)	BTL -1	Understanding	PO2,PO5
7.	State the third order R.K method algorithm to find the numerical solution of the first order differential equation. (A.U M/J 2011,N/D 2012)	BTL -1	Remembering	PO1
8.	Write Milne’s predictor formula and Milne’s corrector formula. (A.U M/J 2012,N/D 2014)	BTL -1	Understanding	PO1
9.	Write down Adams-Bashforth Predictor and Adams-Bashforth corrector formula. (A.U N/D 2011)	BTL -1	Understanding	PO1
10.	State Euler formula. (A.U M/J 2013)	BTL -1	Understanding	PO1
11.	Write down finite difference formula for $y'(x)$ and $y''(x)$ (A.U M/J 2012,N/D 2014)	BTL -1	Understanding	PO1
12.	Write down the Taylor series formula for solving first order ODE.	BTL -1	Understanding	PO1
13.	Using Taylor series method, find the value of $y(0.1)$, from $f(x,y)=x^2 + y^2$ and $y(0) = 1$ correct to 4 decimal places	BTL -4	Analyzing	PO2
14.	Compare Taylor series method and RungeKutta method.	BTL -2	Remembering	PO5
15.	What are the advantages of R-K method over Taylor series method? (A.U N/D 2017)	BTL -2	Remembering	PO5
16.	Compare Single-step method Multi-step methods	BTL -1	Remembering	PO1
17.	Write down the error in Adam’s predictor and corrector formulas	BTL -1	Understanding	PO1
18.	Write down the error in Milne’s predictor and corrector formulas	BTL -1	Understanding	PO1
19.	Compare Adam’s Bashforth method with RungeKutta method	BTL -1	Understanding	PO1
PART-B				
1.	Using Taylor’s series method find y at $x = 0.1$ if $f(x,y)=x^2y - 1, y(0) = 1$	BTL -1	Remembering	PO1,PO2,PO5
2	Solve: $y'' = x + y$; $y(0) = 1$, by Taylor’s series method. Find the values y at $x = 0.1$ and $x = 0.2$	BTL -3	Applying	PO1,PO2,PO5
3	Using Taylor’s series method find $y(1.1)$ given $y'' = x + y, y(1) = 0$	BTL -1	Remembering	PO1,PO2,PO5

4	Using Euler's method find $y(0.2)$ and $y(0.4)$ from $y'' = x + y, y(0) = 1$ with $h = 0.2$	BTL -1	Remembering	PO1,PO2,PO5
5	Consider the initial value problem $y'' = y - x^2 + 1, y(0) = 0.5$ using the modified Euler's method, find $y(0.2)$	BTL -2	Understanding	PO1,PO2,PO5
6	Using R.K method of fourth order, Solve $\frac{dy}{dx} = \frac{y^2 - x^2}{y^2 + x^2}$ with $y(0) = 1$ at $x = 0.2$.	BTL -1	Remembering	PO1,PO2,PO5
7	Using Milne's method find $y(4.4)$ given $5xy'' + y^2 - 2 = 0$ given $y(4) = 1, y(4.1) = 1.0049, y(4.2) = 1.0097$ and $y(4.3) = 1.0143$.	BTL -1	Remembering	PO1,PO2,PO5
8	Obtain the approximate value of y at $x = 0.1$ & 0.2 for the differential equation $\frac{dy}{dx} = 2y + 3e^x, y(0) = 0$ by Taylor's Series method. Compare the numerical solution obtained with the exact solution	BTL -3	Applying	PO1,PO2,PO12
9	Solve $\frac{dy}{dx} = \sin x + \cos y, y(2.5) = 0$ by Modified Euler's method by choosing $h = 0.5$, find $y(3.5)$	BTL -3	Applying	PO1,PO2,PO12
10	Solve $(1+x)\frac{dy}{dx} = -y^2, y(0) = 1$ by Modified Euler's method by choosing $h = 0.1$, find $y(0.1)$ and $y(0.2)$	BTL -3	Applying	PO1,PO2,PO12
11	Apply Runge – Kutta method, to find an approximate value of y when $x = 0.2$ given that $\frac{dy}{dx} = x + y, y(0) = 1$.	BTL -5	Evaluating	PO1,PO2,PO5
12	Given $\frac{dy}{dx} = x - y^2, y(0) = 0, y(0.2) = 0.02, y(0.4) = 0.0795$ and $y(0.6) = 0.1762$. Compute $y(1)$ using Milne's Method.	BTL -3	Applying	PO1,PO2,PO12
13	Using Milne's method to find $y(4.4)$ given that $5xy' + y^2 - 2 = 0$ given that $y(4) = 1, y(4.1) = 1.0049, y(4.2) = 1.0097, y(4.3) = 1.0143$	BTL -1	Remembering	PO1,PO2,PO5

ANSWERS FOR TWO MARK QUESTIONS

UNIT -I-TESTING OF HYPOTHESIS

PART-A(2 MARKS)

<p>(1).What is statistical hypothesis?(Nov/Dec-2017)</p> <p>A statistical hypothesis is a hypothesis concerning the parameters or from of the probability distribution for a designated population or populations, or, more generally, of a probabilistic mechanism which is supposed to generate the observations</p>
<p>(2).Define chi-square.?(Nov/Dec-2017)</p> $\chi^2 = \sum_{i=1}^n (O_i - E_i)^2 / E_i$
<p>(3)Write type I and type II errors.(Apr/May-2015)(Nov/Dec-2013)(May/Jun-2016)</p> <p>Type I error : Rejecting H_0 when is true.</p> <p>Type II error : Accepting H_0 when it is false.</p>
<p>(4) What are the assumptions in 't' distribution?(Nov/Dec-2016)(Apr/May-2015)</p> <p>(i) The parent population from which the sample is drawn is normal.</p> <p>(ii) The sample is random.</p>
<p>(5) State the important properties of the t-distribution.(Apr/May-2015)</p> <p>(i) For sufficiently large value of n,the t-distribution tends to the standard normal distribution.</p> <p>(ii) The mean of the t-distribution is zero</p> <p>(iii). The probability curve of the t-distribution is similar to the std.normal curve and is symmetric about t=0,bell-shaped.</p>
<p>6). Write any three applications of Chi-Square distribution.(May/Jun-2014)</p> <p>(i) To test the goodness of fit.</p> <p>(ii) to test the independence of attributes.</p> <p>(iii)To test the homogeneity of independent estimates of population.</p>
<p>(7) Define null and alternative hypothesis.</p> <p>For applying the tests of significance ,we first set up a hypothesis which is a definite statement about the population parameter called Null hypothesis.Any hypothesis which is complementary to null hypothesis is called an alternative hypothesis.</p>
<p>(8) When do we use the t-distribution?</p> <p>When the sample size is 30 or less and the population standard deviation is unknown,we use the t-distribution.</p>
<p>(9) What is meant by level of significance?</p> <p>The probability 'α' that a random value of the statistic 't' belongs to the critical region is known as level of significance.</p>
<p>(10) Define Standard error and Critical region.</p> <p>The standard deviation of the sampling distribution of a statistic is known as the standard error.</p> <p>A region corresponding to a statistic 't' in the sample S amounts to rejection of the null hypothesis is called critical region.</p>
<p>(11) Write any two applications of 't'-distribution.</p> <p>The t-distribution is used to test the significance of the difference between</p> <p>(i) the mean of the small sample and mean of the population.</p> <p>(ii) The coefficient of correlation in the small sample and that in the population</p>

assumed zero.
(12) Write the condition for the application of χ^2 test. (i) The sample observations should be independent. (ii) N, the total frequency should be at least 50. (iii) Theoretical cell frequency should be less than 5.
(13) Write any three applications of 'F' distribution. F-test is used to test whether (i) Two independent samples have been drawn from the normal populations with the same variance σ^2 . (ii) Two independent estimates of the population variance are homogeneous or not.
(14) State the important properties of F-distribution. (i) The square of the t-variate with n degrees of freedom follows a F-distribution with 1 and n degrees of freedom.
15) Define sampling distribution. Different samples from the same population will result in general in distinct estimates, will form a statistical distribution called sampling distribution.
(16) Define Chi-square test of goodness of fit. Chi-square test of goodness of fit is a test to find if the deviation of the experiment from theory is just by chance or it is due to the inadequacy of the theory to fit the observed data.
(17) Write down the form of the 95% confidence interval for the population mean in terms of population S.D. $\left(\bar{X} - 1.96 \frac{\sigma}{\sqrt{n}}, \bar{X} + 1.96 \frac{\sigma}{\sqrt{n}} \right)$
(18) What is the Standard error of the difference between the means of two large samples drawn from different populations with known SD's. $\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$
(19) What is the test statistic used to test the significance of the difference between small sample mean and population? $t = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$
(20) What is the test statistic used to test the significance of the difference between the means of two small samples? $t = \frac{\bar{x}_1 - \bar{x}_2}{\sigma \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$
(21) Write down the formula of test statistic 'Z' to test the significance of difference between the means (large samples).

$$Z = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

(22) Write down the formula of test statistic 'Z' to test the significance of difference between the proportions (large samples).

$$Z = \frac{p_1 - p_2}{\sqrt{\frac{P_1Q_1}{n_1} + \frac{P_2Q_2}{n_2}}}$$

(23) What is the test statistic used to test the significance of the difference between the means of two small samples of the same size, when the sample items are correlated?

$$t = \frac{\bar{d}}{s\sqrt{n-1}}, \text{ where } d_i = x_i - y_i$$

(24) What are the expected frequency of 2x2 contingency table given below.

a	b
c	d

$\frac{(a+b)(a+c)}{N}$	$\frac{(a+b)(b+d)}{N}$
$\frac{(a+c)(d+c)}{N}$	$\frac{(d+b)(d+c)}{N}$

(25) Write down the 1% and 5% critical values for right tailed and two tailed Tests.

	1%	5%
Two tailed test :	2.58	1.96
Right tailed test :	2.33	1.645

(26) What is the difference between confidence limits and tolerance limits

Confidence limits: To estimate a parameter of a population

Tolerance limits: To indicate between what limits one can find a certain proportion of a population.

(27) What are the assumptions of large sample?

(i) it should be normal

(ii) values given by the samples are sufficiently close to the population parameters.

(28) What is test of goodness of fit?

To determine whether the actual sample distribution matches a known theoretical distribution.

(29) Define hypothesis

Hypothesis is a statement about the population parameter. It is tested on the basis of the outcome of the random sample.

There are 2 types (i) null hypothesis and (ii) alternate hypothesis

(30) What is meant by population?

A population in statistics means a set of objects which are measurement or observations pertaining to the objects.

**UNIT -II-DESIGN OF EXPERIMENTS
PART-A(2 MARKS)**

1).Write the advantages of Latin Square design.(Nov/Dec-2017)

Advantages of latin square designs. Controls more variation than CR or RCB designs because of 2-way stratification.

(2).What are the conditions to be followed in one way classification?(Nov/Dec-2017)

In statistics, **one-way analysis of variance** (abbreviated **one-way ANOVA**) is a technique that can be used to compare means of two or more samples (using the F distribution). This technique can be used only for numerical response data, the "Y", usually one variable, and numerical or (usually) categorical input data, the "X", always one variable, hence "one-way"

(3).What is meant by analysis of variance?(May/Jun-2016)

Analysis of Variance is a technique that will enable us to test for the significance of the difference among more than two sample means.

4).Why a 2x2 Latin square is not possible?Explain.(May/Jun-2016)(May/Jun-2014).

Consider a nxn latin Square design ,then the degrees of freedom for SSE

$$= (n^2 - 1) - (n - 1) - (n - 1) - (n - 1)$$

$$= (n - 1)(n - 2)$$

For n=2 ,degrees of freedom of SSE=0 and hence MSE id not defined.Comparision is not possible.Hence 2x2 Latin Square is not possible.

(5)Define Replication and Randomization.(Nov/Dec-2016)

Replication is the repetition of an experimental condition so that the variability associated with the phenomenon can be estimated. In other words replication as "the repetition of the set of all the treatment combinations to be compared in an experiment. Each of the repetitions is called a **replicate**."

A method based on chance alone by which study participants are assigned to a treatment group. **Randomization** minimizes the differences among groups by equally distributing people with particular characteristics among all the trial arms.

(6) What is the advantage of factorial experiment?(Nov/Dec-2016)

(i)Factorial designs allow additional factors to be examined at no additional cost

(ii)Factorial designs allow the effects of a factor to be estimated at several levels of the other factors, yielding conclusions that are valid over a range of experimental conditions.

(7)What is the aim of design of experiment?(Apr/May-2015)(May/Jun-2014)

The **design of experiments** (DOE, DOX, or **experimental design**) is the **design** of any task that **aims** to describe or explain the variation of information under conditions that are hypothesized to reflect the variation.

(8) What are the basic principles of experimental design?(Apr/May-2015)

(i) Replication

(ii) Randomization and Local control.
(9) Write the advantages and disadvantages of RBD?(Apr/May-2015) Advantages : (i). Accuracy (ii) Flexibility (iii) Easy to analyze Disadvantage : It is not suitable for large number of treatment
(10) What is Latin Square design ? A useful method of eliminating fertility variations consist in an experimental layout which will control in 2 perpendicular directions such a layout is a LSD.
(11) Define Raw Sum of Squares and Correction factor. The expression $\sum \sum x_{ij}^2$ is known as RSS and the expression $\frac{G^2}{N}$, where $G^2 = \sum \sum x_{ij}^2$ is called the correction factor.
(12) Write any 3 applications of LSD. (i) The statistical analysis is simple. (ii) Even with the missing data analysis remains relatively simple. (iii) More than one factor can be investigated simultaneously.
(13) How do you calculate the Correction factor in LSD? By squaring the grand total and dividing it by the number of observations ,we calculate the correction factor.
(14) What do you mean by design of experiments?(Nov/Dec-2014) It is defined as the logical construction of the experiment in which the degree of uncertainty with which the inference is drawn ,may be well defined.
(15) What are the subject matters included in the design of experiment? (i) Planning of the experiment. (ii) Obtaining relevant information from it regarding the statistical hypothesis under study.
(16) What are the assumptions in ANOVA? Each of samples is drawn from a normal population and the variances for the population from which samples have been drawn are equal.
(17) What are the three essential steps to plan an experiment? (i) A statement of the objective. (ii) Statement should clearly mention the hypothesis to be tested. (iii) Description should include the type of experimental material, size of the experiment and the number of replications.
(18) What are the basic steps in ANOVA? (i) Estimate the population variance among the sample means. (ii) Estimate the population variance from the variance within the sample means.
(19) Write the steps to find F-ratio. $F = \frac{S_1^2}{S_2^2} = \frac{\text{Variance between samples}}{\text{Variance within samples}}$

<p>(20) Discuss the advantages of Completely Randomized block design.</p> <p>(i) easy to lay out (ii) allows flexibility (iii) simple statistical information (iv). The lot of information due to missing data is smaller than with any other design</p>
<p>(21) State the uses of ANOVA.</p> <p>(i) The effects of some fertilizer on the yields are significantly different. (ii) The mean qualities of outputs of various machines differ significantly.</p>
<p>(22) Explain the word treatment in ANOVA.</p> <p>The word treatment in ANOVA is used to refer to any factor in experiment is controlled at different levels or values.</p>
<p>(23) What do you mean by 2-way classification?</p> <p>In two way classification ,the datas are classified according to different criteria or factors.</p>
<p>(24) Indicate the characteristics of a good experimental design.</p> <p>(i) Absolute (ii) Comparative.</p>
<p>(25) What are the important designs of experiments?</p> <p>(i) Completely Randomized design (or) One-Way classification (ii) Randomized Block Design (or) Two-Way classification (iii) Latin Square Design (or) Three-Way classification.</p>
<p>(26) What is an experimental error ?</p> <p>The variation from plot to plot caused by uncontrolled factors is known as experimental error.</p>
<p>(27) What is meant by CRD?</p> <p>It is defined as a type of experimental design where the experimental units are allocated to the treatments in a completely random fashion. This is used to study the effects of one primary factor without the need to take other nuisance variables into account.</p>
<p>(28) Compare RBD and LSD.</p> <p>RBD is more efficient than CRD for most types of experiment work. In CRD, grouping of the experiments size so as to allocate the treatments at random to the experimental units is not done. But in RBD ,treatments are allocated at random within the units of each stratum. RBD is more flexible than CRD, since no restrictions are placed on the number or treatments or the number of replicates.</p>
<p>(29) Compare LSD and RBD.</p> <p>In LSD, the number of treatments is equal to the number of replications ,whereas there are no such restrictions on treatments and replications in RBD.</p>
<p>(30) What are the uses of Chi-Square test?</p> <p>(i) To test significance difference between experimental values and theoretical values. (ii) To find whether two or more attributes are associated or not.</p>

UNIT III SOLUTION OF EQUATIONS AND EIGENVALUE PROBLEMS

Newton Raphson method – Gauss elimination method – pivoting – Gauss Jordan methods – Iterative

methods of Gauss Jacobi and Gauss Seidel – Matrix inversion by Gauss Jordan method – Eigen values of a matrix by power method.

Textbook : Grewal. B.S., and Grewal. J.S., "Numerical Methods in Engineering and Science", 9th Edition, Khanna Publishers, New Delhi, 2007.

1.State the order (rate) of convergence and convergence condition for Newton Raphson method.

Sol. The order of convergence of Newton Raphson method is 2

(quadratic) and convergence condition is $|f(x)f''(x)| < [f'(x)]^2$.

2. Give Newton Raphson iterative formula.

Sol. $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}, n = 0, 1, 2, \dots$

3. Establish an iteration formula to find the reciprocal of a positive number N by Newton Raphson method.

Sol. Let $x = 1/N$

$$\Rightarrow N = \frac{1}{x} \Rightarrow \frac{1}{x} - N = 0$$

$$(i.e.) f(x) = \frac{1}{x} - N \Rightarrow f(x_n) = \frac{1}{x_n} - N, f'(x_n) = -\frac{1}{x_n^2}$$

By Newton Raphson method,

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{\frac{1}{x_n} - N}{-\frac{1}{x_n^2}} = x_n + x_n^2 \left(\frac{1}{x_n} - N \right) = x_n (2 - Nx_n).$$

4. State the principle used in Gauss-Jordan method.

Sol. In the equation $AX = B$, the matrix A is transformed into an identity matrix.

5. Give the sufficient condition of convergence of Gauss Seidel method.

Sol. The absolute value of the leading diagonal element is greater than the sum of the absolute values of the other elements in that row, which is called diagonally dominant.

6. Write the conditions for convergence in Gauss Seidel iterative technique. (or) When the method of iteration will be useful?

Sol. The coefficient matrix should be diagonally dominant.

7. State Gauss Seidel method.

Sol. As soon as a new value for a variable is found by iteration it is used immediately in the following equations. This method is called Gauss Seidel method.

8. Gauss Seidel method always converges – True or False.

Sol. False.

9. Write the first iteration values of x,y,z when the equations $27x+6y-z = 85$, $6x+15y+2z = 72$, $x+y+5z = 110$ are solved by Gauss Seidel method.

Sol. Here the coefficient matrix is diagonally dominant. Then

$$x = \frac{1}{27}(85 - 6y + z) \dots\dots(1)$$

$$y = \frac{1}{15}(72 - 6x - 2z) \dots\dots(2)$$

$$z = \frac{1}{5}(110 - x - y) \dots\dots(3)$$

First Iteration

Put $y = 0, z = 0$ in (1), we get $x = 3.148$

Put $x = 3.148, z = 0$ in (2), we get $y = 3.451$

Put $x = 3.148, y = 3.451$ in (3), we get $z = 20.662$

10. Compare Gauss Elimination and Gauss Jordan methods for solving linear systems of the form $AX=B$.

Sol. In Gauss Elimination method, the coefficient matrix reduced to upper triangular matrix and we get the solution by back substitution whereas in Gauss Jordan method, the coefficient matrix reduces to an unit or identity matrix and we get the solution without using back substitution.

11. What type of Eigen value can be obtained using power method?

Sol. Dominant eigen value.

12. Find the dominant eigen value of $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ by power method.

Sol. Dominant eigen value = 5.3722

13. On what type of equations Newton's method can be applicable ?

Sol. Newton's method can be applicable to the solution of both algebraic and transcendental equation and can be also used when the roots are complex.

14. By Gauss elimination method solve $x + y = 2$ and $2x + 3y = 5$.

Sol. The augmented matrix is

$$[A, B] = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \end{bmatrix} R_2 = R_2 - 2R_1$$

By back substitution, $x + y = 2$ ----(1)

$$y = 1$$

(1) becomes, $x + 1 = 2$

$$x = 1$$

Hence $x = 1, y = 1$.

15. Why Gauss Seidel iteration is a method of successive corrections?

Sol. Because we replace approximations by corresponding new ones as soon the latter have been computed.

16. What are the merits of Newton's method of iteration?

Sol. Newton's method is successfully used to improve the result obtained by other methods. It is applicable to the solution of equations involving algebraical functions as well as transcendental functions.

17. Give two direct methods to solve a system of linear equations.

Sol. Gauss Elimination method and Gauss Jordan method.

18. Compare Gauss Elimination with Gauss Seidel method.

Sol. Gauss Elimination	Gauss Seidel
i. Direct method	i. Indirect method
ii. Used to find inverse of the matrix also.	ii. Used to solve system of equations only
iii. Diagonally dominant condition is not insisted.	iii. Diagonally dominant condition is insisted.

UNIT IV INTERPOLATION, NUMERICAL DIFFERENTIATION AND NUMERICAL INTEGRATION

Lagrange's and Newton's divided difference interpolations – Newton's forward and backward difference interpolation – Approximation of derivatives using interpolation polynomials – Numerical single and double integrations using Trapezoidal and Simpson's 1/3 rules.
 Textbook : Grewal. B.S., and Grewal. J.S., "Numerical Methods in Engineering and Science", 9th Edition, Khanna Publishers, New Delhi, 2007.

1. Define interpolation and extrapolation?

Sol. The process of computing the value of a function inside the given range is called interpolation. The process of computing the value of a function outside the given range is called extrapolation.

2. State Newton's formula on interpolation. When it is used?

Sol. Newton's forward interpolation formula is

$$y = y_0 + u\Delta y_0 + \frac{u(u-1)}{2!} \Delta^2 y_0 + \frac{u(u-1)(u-2)}{3!} \Delta^3 y_0 + \dots$$

$$\text{where } u = \frac{x - x_0}{h}$$

This formula is used mainly for interpolating the values of y near the beginning of a set of tabular values.

Newton's backward interpolation formula is

$$y = y_n + u\nabla y_n + \frac{u(u+1)}{2!} \nabla^2 y_n + \frac{u(u+1)(u+2)}{3!} \nabla^3 y_n + \dots$$

$$\text{where } u = \frac{x - x_n}{h}$$

This formula is used mainly for interpolating the values of y near the end of a set of tabular values.

3. Say True or False. – Newton's divided difference formula is applicable only for equally spaced intervals.

Sol. False.

4. State Newton's divided difference formula.

Sol. $y = y_0 + (x - x_0)\Delta y_0 + (x - x_0)(x - x_1)\Delta^2 y_0 + (x - x_0)(x - x_1)(x - x_2)\Delta^3 y_0 + \dots$

5. State Lagrange's interpolation formula

Sol.

$$y = f(x) = \frac{(x - x_1)(x - x_2)(x - x_3) \dots (x - x_n)}{(x_0 - x_1)(x_0 - x_2)(x_0 - x_3) \dots (x_0 - x_n)} y_0$$

$$+ \frac{(x - x_0)(x - x_2)(x - x_3) \dots (x - x_n)}{(x_1 - x_0)(x_1 - x_2)(x_1 - x_3) \dots (x_1 - x_n)} y_1$$

$$+ \frac{(x - x_0)(x - x_1)(x - x_3) \dots (x - x_n)}{(x_2 - x_0)(x_2 - x_1)(x_2 - x_3) \dots (x_2 - x_n)} y_2$$

$$+ \dots +$$

$$+ \frac{(x - x_0)(x - x_1)(x - x_2)(x - x_3) \dots (x - x_{n-1})}{(x_n - x_0)(x_n - x_1)(x_n - x_2)(x_n - x_3) \dots (x_n - x_{n-1})} y_n$$

6. Use Lagrange's formula to find the quadratic polynomial that takes these values

$$x : 0 \quad 1 \quad 3$$

$$y : 0 \quad 1 \quad 0$$

Then find y(2).

Sol. By Lagrange's formula

$$y = f(x) = \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)} y_0 + \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)} y_1$$

$$+ \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)} y_2$$

$$y = f(x) = \frac{(x - 1)(x - 3)}{(0 - 1)(0 - 3)} \cdot 0 + \frac{(x - 0)(x - 3)}{(1 - 0)(1 - 3)} \cdot 1 + \frac{(x - 0)(x - 1)}{(3 - 0)(3 - 1)} \cdot 0$$

$$y(x) = \frac{x^2 - 3x}{-2}$$

Hence y(2) = 1.

7. By differentiating Newton forward and backward difference formula, find the first derivative of the function f(x).

Sol. Newton forward interpolation formula is

$$y = y_0 + u\Delta y_0 + \frac{u(u-1)}{2!} \Delta^2 y_0 + \frac{u(u-1)(u-2)}{3!} \Delta^3 y_0 + \dots$$

$$\text{where } u = \frac{x - x_0}{h}$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

$$= \frac{1}{h} \left[\Delta y_0 + \frac{2u-1}{2} \Delta^2 y_0 + \frac{3u^2 - 6u + 2}{6} \Delta^3 y_0 \right]$$

$$+ \frac{2u^3 - 9u^2 + 11u - 3}{12} \Delta^4 y_0 + \dots]$$

Newton backward interpolation formula is

$$y = y_n + u \nabla y_n + \frac{u(u+1)}{2!} \nabla^2 y_n + \frac{u(u+1)(u+2)}{3!} \nabla^3 y_n + \dots$$

$$\text{where } u = \frac{x - x_n}{h}$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

$$= \frac{1}{h} \left[\nabla y_n + \frac{2u+1}{2} \nabla^2 y_n + \frac{3u^2 + 6u + 2}{6} \nabla^3 y_n \right.$$

$$\left. + \frac{2u^3 + 9u^2 + 11u + 3}{12} \nabla^4 y_n + \dots \right]$$

8. Write down the Newton – cotes quadrature formula..

Sol.

$$\int_{x_0}^{x_n} f(x) dx = h \left\{ n y_0 + \frac{n^2}{2} \Delta y_0 + \frac{1}{2} \left(\frac{n^3}{3} - \frac{n^2}{2} \right) \Delta^2 y_0 + \frac{1}{6} \left(\frac{n^4}{4} - n^3 + n^2 \right) \Delta^3 y_0 + \dots \right\}$$

9. What is the geometrical interpretation of Trapezoidal rule?

Sol. We are finding the area of the curve enclosed by $y = f(x)$, the X-axis, the ordinates $x = a$ and $x = b$ by using the area of trapezium.

10. Using Trapezoidal rule evaluate $\int_0^\pi \sin x dx$ by dividing the range into 6 equal parts.

Sol.
$$h = \frac{\pi - 0}{6} = \frac{\pi}{6}$$

When $h = \frac{\pi}{6}$, the values of $y = \sin x$ are

x :	0	$\frac{\pi}{6}$	$\frac{2\pi}{6}$	$\frac{3\pi}{6}$	$\frac{4\pi}{6}$	$\frac{5\pi}{6}$	π
y=sinx:	0	0.5	.8660	1	.8660	0.5	0

Trapezoidal rule is

$$\int_0^\pi \sin x dx = \frac{h}{2} [(y_0 + y_n) + 2(y_1 + y_2 + y_3 + \dots + y_{n-1})]$$

$$= \frac{\pi}{6(2)} [(0 + 0) + 2(0.5 + 0.8660 + 1 + 0.8660 + 0.5)]$$

$$= 0.9770$$

11. Why is Trapezoidal rule so called?

Sol. The Trapezoidal rule is so called, because it approximates the integral by the sum of n

trapezoids.

12. What are the truncation errors in Trapezoidal and Simpson's rules of numerical integration?

Sol. Error in the Trapezoidal rule is $-\frac{h^3}{12} f''(\theta)$. Error in the Trapezoidal rule is of the order h^2 .

Error in the Simpson's one-third rule is $-\frac{h^5}{90} f^{IV}(\theta)$. Error in Simpson's one-third rule is of the order h^4 .

Error in the Simpson's three eighth rule is $-\frac{3h^5}{80} f^{IV}(\theta)$. Error in the Simpson's three eighth rule is of the order h^4 .

13. What is the condition for Simpson's 3/8 rule and state the formula.

Sol. The condition for Simpson's 3/8 rule is the number of sub-intervals should be a multiple of 3. Simpson's 3/8 rule is

$$\int_{x_0}^{x_n} f(x) dx = \frac{3h}{8} [(y_0 + y_n) + 3(y_1 + y_2 + y_4 + y_5 + y_7 + \dots) + 2(y_3 + y_6 + y_9 + \dots)]$$

14. Using Simpson's rule find $\int_0^4 e^x dx$ given $e^0 = 1, e^1 = 2.72, e^2 = 7.39, e^3 = 20.09, e^4 = 54.6$

Soln The following data is

x :	0	1	2	3	4
y :	1	2.72	7.39	20.09	54.6

Simpson's 1/3rd rule is

$$\int_{x_0}^{x_n} f(x) dx = \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + y_5 + \dots) + 2(y_2 + y_4 + y_6 + \dots)]$$

$$\int_0^4 e^x dx = \frac{1}{3} [(1 + 54.6) + 4(2.72 + 20.09) + 2(7.39)]$$

$$= 53.8733$$

15. Compare Trapezoidal rule and Simpson's 1/3rd rule for evaluating numerical integration.

Sol. i) In Newton Cotes Quadrature formula, if we put $n = 1$ we get

Trapezoidal rule whereas if we put $n = 2$, we get Simpson's 1/3rd rule.

ii) In Trapezoidal rule, the interpolating polynomial is linear whereas in Simpson's 1/3rd rule, the interpolating polynomial is of degree 2.

iii) In Trapezoidal rule, there is no restriction on the number of intervals whereas in Simpson's 1/3rd rule, the number of intervals should be even.

UNIT V NUMERICAL SOLUTION OF ORDINARY DIFFERENTIAL EQUATIONS

Taylor's series method - Euler's method - Modified Euler's method - Fourth order Runge-Kutta method for solving first and second order equations - Milne's predictor-corrector methods for solving first order equations - Finite difference methods for solving second order equation.

1.State Modified Euler algorithm to solve $y' = f(x, y), y(x_0) = y_0$ at $x=x_0 +h$. (A.U.N/D 2017, N/D 2011,2012, M/J 2013)

$$y_{n+1} = y_n + hf \left[x_n + \frac{h}{2}y_n + \frac{h}{2}f(x_n, y_n) \right]$$

$$y_1 = y_0 + hf \left[x_0 + \frac{h}{2}y_0 + \frac{h}{2}f(x_0, y_0) \right]$$

2. State the disadvantage of Taylor series method.

(A.U N/D 2009,M/J 2012,2014)

Solution:

In the differential equation $f(x, y), \frac{dy}{dx} = f(x, y)$ the function $f(x, y)$, may have a complicated algebraical structure. Then the evaluation of higher order derivatives may become tedious. This is the demerit of this method.

3. Write the merits and demerits of the Taylor method of solution. (A.U.N/D 2010, M/J 2012)

Solution:

The method gives a straight forward adaptation of classic to develop the solution as an infinite series. It is a powerful single step method if we are able to find the successive derivatives easily. If $f(x,y)$ involves some complicated algebraic structures then the calculation of higher derivatives becomes tedious and the method fails. This is the major drawback of this method. However the method will be very useful for finding the starting values for powerful methods like Runge - Kutta method, Milne's method etc.,

4.Which is better Taylor's method or R. K. Method?(or) State the special advantage of Runge-Kutta method over taylor series method (A.U M/J 2011)

Solution:

- R.K Methods do not require prior calculation of higher derivatives of $y(x)$, as the Taylor method does. Since the differential equations using in applications are often complicated, the calculation of derivatives may be difficult.
- Also the R.K formulas involve the computation of $f(x, y)$ at various positions, instead of derivatives and this function occurs in the given equation.

5. Compare Runge-Kutta methods and predictor – corrector methods for solution of initial value problem. (A.U M/J 2011)

Solution:

Runge-Kutta methods

- Runge-methods are self starting, since they do not use information from previously calculated points.
- As these are self starting, an easy change in the step size can be made at any stage. Since these methods require several evaluations of the function $f(x, y)$, they are time consuming.
- In these methods, it is not possible to get any information about truncation error.

Predictor Corrector methods

- These methods require information about prior points and so they are not self starting.
- In these methods it is not possible to get easily a good estimate of the truncation error.

6. What is a Predictor-corrector method of solving a differential equation? (A.U M/J 2009)

Solution:

- Predictor-corrector methods are methods which require the values of y at $x_n, x_{n-1}, x_{n-2}, \dots$ for computing the value of y at x_{n+1}
- We first use a formula to find the value of y at x_{n+1} and this is known as a predictor formula.

The value of y so got is improved or corrected by another formula known as corrector formula

7. State the third order R.K method algorithm to find the numerical solution of the first order differential equation. (A.U M/J 2011, N/D 2012)

Solution: To solve the differential equation $y' = f(x, y)$ by the third order R.K method, we use the following algorithm.

$$\begin{aligned}
 k_1 &= hf(x, y) \\
 k_2 &= hf\left(x + \frac{h}{2}, y + \frac{k_1}{2}\right) \\
 k_3 &= hf(x + h, y + 2k_2 - k_1) \\
 \text{and } \Delta y &= \frac{1}{6}(k_1 + 4k_2 + k_3)
 \end{aligned}$$

8. Write Milne's predictor formula and Milne's corrector formula.

(A.U M/J 2012, N/D 2014)

Solution:

- Milne's predictor formula is

$$y_{4,p} = y_0 + \frac{4h}{3} [2y'_1 - y'_2 + 2y'_3]$$

- Milne's corrector formula is

$$y_{4,c} = y_2 + \frac{h}{3} [y'_2 + 4y'_3 + y'_4]$$

9. Write down Adams-Bashforth Predictor and Adams-Bashforth corrector formula.

(A.U N/D 2011)

<p>Solution Adams-Bashforth predictor formula is</p> $y_{4,p} = y_3 + \frac{h}{24} [55y'_3 - 59y'_2 + 37y'_1 - 9y'_0]$ <p>Adams-Bashforth corrector formula is</p> $y_{4,c} = y_3 + \frac{h}{24} [9y'_4 + 19y'_3 - 5y'_2 + y'_1]$								
<p>10. State Euler formula (A.U M/J 2013)</p> <p>Solution: $y_{n+1} = y_n + hf[x_n, y_n]$ when $n = 0, 1, 2, \dots$</p>								
<p>11. Write down finite difference formula for $y'(x)$ and $y''(x)$ (A.U M/J 2012, N/D 2014)</p> <p>Solution:</p> $y'(x) = \frac{y_{i+1} - y_i}{h}, \quad y''(x) = \frac{y_{i-1} - 2y_i + y_{i+1}}{h^2}$								
<p>12. Write down the Taylor series formula for solving first order ODE.</p> $y_{n+1} = y_n + \frac{h}{1!} y'_n + \frac{h^2}{2!} y''_n + \frac{h^3}{3!} y'''_n + \dots$								
<p>13. Using Taylor series method, find the value of $y(0.1)$, from $dy = x^2 + y^2$ and $y(0) = 1$ correct to 4 decimal places</p> <p>Solution:</p> $y' = x^2 + y^2 \quad \begin{matrix} x=0 \\ y=1 \end{matrix}$ $y'_0 = 2x + 2yy' \quad \begin{matrix} y'_0 = 2 \\ y''_0 = 8 \\ y'''_0 = 28 \end{matrix}$ <p>By using Taylor series formula, $y_1 = 1.11145$</p>								
<p>14. Compare Taylor series method and Runge Kutta method.</p> <p>Solution:</p> <ul style="list-style-type: none"> ➤ The use of R-K method gives quick convergence to the solutions of the differential equations than Taylor's series method. ➤ The labour involved in R-K method is comparatively lesser. ➤ In R-K method, the derivatives of higher order are not required for calculation as in Taylor series method. 								
<p>15. What are the advantages of R-K method over Taylor series method?</p> <p>Solution: The Rungekutta methods are designed to give greater accuracy and they possess the advantage of requiring only the function values at some selected points on the sub interval.</p>								
<p>16. Compare Single-step method Multi-step methods. (A.U N/D 2017)</p> <p>Solution:</p> <table border="1" style="width: 100%; border-collapse: collapse; margin-top: 10px;"> <thead> <tr> <th style="width: 15%;">S.No</th> <th style="width: 40%;">Single-step method</th> <th style="width: 45%;">Multi-step method</th> </tr> </thead> <tbody> <tr> <td> </td> <td> </td> <td> </td> </tr> </tbody> </table>			S.No	Single-step method	Multi-step method			
S.No	Single-step method	Multi-step method						

1	It requires only the numerical value y_i in order to compute the next value y_{i+1}	It requires not only the numerical value y_i but also at least four of the past values y_{i-1}, y_{i-2}, \dots
2	Taylor series, Euler's and R-K methods are single step methods	Milne's, Adam's methods are multi step methods

17. Write down the error in Adam's predictor and corrector formulas.

Solution:

Order of error is h^5

Error in predictor $h^5 \frac{251}{720} y^{(v)}$

Error in corrector $h^5 \frac{19}{120} y^{(v)} \xi$

18. Write down the error in Milne's predictor and corrector formulas.

Solution:

Order of error is h^5

Error in predictor $h^5 \frac{14}{45} y^{(v)} \xi$

Error in corrector $h^5 \frac{1}{90} y^{(v)} \xi$

19. Compare Adam's Bashforth method with RungeKutta method

Solution:

S. No	Adam's Bashforth Method	Runge-Kutta Method
1	Multi step method	Single step method
2	Need four prior values of y_i 's	Need only the last prior value
3	Does not permits changes in the step size	Permits changes in the step size